

COSMIC LITHIUM-BERYLLIUM-BORON STORY

ELISABETH VANGIONI-FLAM

*Institut d'Astrophysique de Paris, CNRS
98 bis bd Arago Paris France*

AND

MICHEL CASSÉ

*Service d'Astrophysique, CEA
Orme des Merisiers
91191 Gif/Yvette France
and Institut d'Astrophysique de Paris*

Abstract.

Light element nucleosynthesis is an important chapter of nuclear astrophysics. Specifically, the rare and fragile light nuclei Lithium, Beryllium and Boron (LiBeB) are not generated in the normal course of stellar nucleosynthesis (except ${}^7\text{Li}$) and are, in fact, destroyed in stellar interiors. This characteristic is reflected in the low abundance of these simple species. Up to recently, the most plausible interpretation was that Galactic Cosmic Rays (GCR) interact with interstellar CNO to form LiBeB. Other origins have been also identified: primordial and stellar (${}^7\text{Li}$) and supernova neutrino spallation (${}^7\text{Li}$ and ${}^{11}\text{B}$). In contrast, ${}^9\text{Be}$, ${}^{10}\text{B}$ and ${}^6\text{Li}$ are pure spallative products. This last isotope presents a special interest since the ${}^6\text{Li}/{}^7\text{Li}$ ratio has been measured recently in a few halo stars offering a new constraint on the early galactic evolution of light elements. Optical measurements of the beryllium and boron abundances in halo stars have been achieved by the 10 meter KECK telescope and the Hubble Space Telescope. These observations indicate a quasi linear correlation between Be and B vs Fe, at least at low metallicity, which, at first sight, is contradictory to a dominating GCR origin of the light elements which predicts a quadratic relationship. As a consequence, the theory of the origin and evolution of LiBeB nuclei has to be refined. Aside GCRs, which are accelerated in the general interstellar medium (ISM) and create LiBeB through the break up of CNO by fast protons and alphas, Wolf-Rayet stars (WR) and core collapse supernovae (SNII) grouped in superbubbles could produce copious amounts of light

elements via the fragmentation in flight of rapid carbon and oxygen nuclei colliding with H and He in the ISM. In this case, LiBeB would be produced independently of the interstellar medium chemical composition and thus a primary origin is expected. These different processes are discussed in the framework of a galactic evolutionary model. More spectroscopic observations (specifically of O, Fe, Li, Be, B) in halo stars are required for a better understanding of the relative contribution of the various mechanisms. Future tests on the injection and acceleration of nuclei by supernovae and Wolf Rayet relying on gamma-ray line astronomy will be invoked in the perspective of the European INTEGRAL satellite.

1. Introduction

A general trend in nature is that complex nuclei are not proliferating: the abundance of the elements versus the mass number draws a globally decreasing curve. In the whole nuclear realm, LiBeB are exceptional since they are both simple and rare. Typically, in the Solar System, $\text{Li}/\text{H} = 2 \cdot 10^{-9}$, $\text{B}/\text{H} = 7 \cdot 10^{-10}$, $\text{Be}/\text{H} = 2.5 \cdot 10^{-11}$ (Anders and Grevesse 1989). Indeed, they are rare because they are fragile and apparently a selection principle at the nuclear level has operated in nature. Due to the fact that nuclei with mass 5 and 8 are unstable, the Big-Bang nucleosynthesis (BBN) has stopped at $A = 7$, and primordial thermonuclear fusion has been unable to proceed beyond lithium. The standard BBN is hopelessly ineffective in generating ${}^6\text{Li}$, ${}^9\text{Be}$, ${}^{10}\text{B}$, ${}^{11}\text{B}$ (fig 1). Thus, stars were necessary to pursue the nuclear evolution bridging the gap between ${}^4\text{He}$ and ${}^{12}\text{C}$ much later, through nuclear fusion.

Stellar nucleosynthesis, quiescent or explosive, forge the whole variety of nuclei from C to U but they destroy LiBeB in the interior of stars, except ${}^7\text{Li}$ which is produced in AGB and novae. The destruction temperature are 2, 2.5, 3.5 and 5.3 millions of degrees for ${}^6\text{Li}$, ${}^7\text{Li}$, ${}^9\text{Be}$, and ${}^{10}\text{B}$ respectively. Finally, ${}^7\text{Li}$ and ${}^{11}\text{B}$ could be produced by neutrino spallation in carbon shells of core collapse supernovae (Woosley *et al.* 1990, Vangioni-Flam *et al.* 1996); however, this mechanism is particularly uncertain depending strongly on the neutrino energy distribution.

It is clear that another source is necessary to generate at least ${}^6\text{Li}$, ${}^9\text{Be}$, ${}^{10}\text{B}$ and this non thermal mechanism is the break up of heavier species (CNO, mainly) by energetic collisions, also called spallation.

The LiBeB story has been rich and moving. The genesis of LiBeB was so obscure to Burbidge *et al.* (1957) that they called X the process leading to their production. Then came Hubert Reeves and his students. In a semi-




Figure 1. Big Bang Nucleosynthesis of Lithium, Beryllium and Boron vs photon over baryon ratio

nal work, Meneguzzi, Audouze and Reeves (1971) identified the production process, i.e. the Galactic Cosmic Rays - Interstellar Medium interaction. Exploiting the fast p, α in the GCRs interacting with CNO in the ISM, they were able to make quantitative estimates of the LiBeB production on the basis of cross section measurements notably made in Orsay (Raisbeck and Yiou, 1971, 1975). However, this estimate, based on the local and present observations (LiBeB and CNO abundances, cosmic ray flux and spectrum) was based on an extrapolation over the whole galactic lifetime assuming that all the parameters are constant. This result accounted fairly well for the cumulated light element abundances but obviously not for their evolution which, at that time, was unknown. The pertinence of their idea is illuminated by the simple and beautiful fact that the hierarchy of the abundances $^{11}\text{B} > ^{10}\text{B} > ^9\text{Be}$ is reflected in the cross sections (Read and Viola, 1984). This is another proof that nature follows the rules of nuclear physics. ^6Li , ^9Be and ^{10}B were nicely explained but problems were encountered with ^7Li and ^{11}B . The calculated $^7\text{Li}/^6\text{Li}$ ratio was 1.2 against 12.5 in meteorites. Stellar sources of ^7Li appeared necessary. The estimated $^{11}\text{B}/^{10}\text{B}$ ratio was 2.5 instead of 4 in meteorites. An ad-hoc hypothesis drawing on unobservable low energy proton operating through the $^{14}\text{N}(p, x)^{11}\text{B}$ reaction was advocated (see Reeves 1994 for a review).

New measurements of Be/H and B/H from KECK and HST, together with [Fe/H] (Rebolo *et al.* 1988, Gilmore *et al.* 1992, Duncan *et al.* 1992, Boesgaard and King 1993, Ryan *et al.* 1994, Duncan *et al.* 1997, García-López *et al.* 1998) in very low metallicity halo stars came to set strong constraints on the origin and evolution of light isotopes.

The evolution of BeB was suddenly known over about 10 Gyr, taking

[Fe/H] as an evolutionary index. A compilation of Be and B data is presented in fig 2. The most striking point is that $\log(\text{Be}/\text{H})$ and $\log(\text{B}/\text{H})$ are both quasi proportional to [Fe/H], at least up to [Fe/H] = -1 and that the B/Be ratio lies in the range 10 - 30 (Duncan *et al.* 1997).




Figure 2. Beryllium and Boron evolution vs [Fe/H]

This linearity came as a surprise since a quadratic relation was expected from the GCR mechanism. It was a strong indication that the standard GCRs are not the main producers of LiBeB in the early Galaxy. A new mechanism of primary nature was required to reproduce these observations: low energy fast CO nuclei produced and accelerated by massive stars (WR and SN II) fragment on H and He at rest in the ISM. This low energy component (LEC) has the advantage of coproducing Be and B in good agreement with the ratio observed in Pop II stars, (figure 2 and Vangioni-Flam *et al.* 1998a).

A primary origin, in this language, means a production rate independent of the interstellar metallicity. In this case, the cumulated abundance of a given light isotope L is approximately proportional to Z. At variance, standard GCRs offer a secondary mechanism because it should depend both on the CNO abundance of the ISM at a given time and on the intensity of cosmic ray flux, itself assumed to be proportional to the SN II rate.

Note however, the two discrepant points in the boron diagram at the lowest [Fe/H]. This is mainly due to the huge NLTE correction on the data (Kiselman and Carlsson 1996) that increases the departure from a straight line. It is important to take a carefull look to this delicate correction.

The Be-Fe and B-Fe correlations taken at face value show a contradiction between theory and observation. But, since oxygen is the main progenitor of BeB, the apparent linear relation between BeB and Fe could be misleading if O were not strictly proportional to Fe (Israelian *et al.* 1998

and Boesgaard *et al.* 1998). Thus the pure primary origin of BeB could be questioned (Fields and Olive 1998). However, the oxygen measurements themselves are confronted to many difficulties (Mac Williams 1997, Cayrel, Spite and Spite, private communication). On the theoretical side, the situation is not better. The $[\alpha/\text{Fe}]$ vs $[\text{Fe}/\text{H}]$ where $\alpha = \text{Mg, Si, Ca, S, Ti}$ (Cayrel 1996, Ryan *et al.* 1996) show a plateau from about $[\text{Fe}/\text{H}] = -4$ to -1 . On nucleosynthetic grounds, it would be surprising that oxygen does not follow the Si and Ca trends. Moreover, using the published nucleosynthetic yields (Woosley and Weaver 1995, Thielemann, Nomoto and Hashimoto 1996) it is impossible to fit the $\log(\text{O}/\text{H})$ vs $[\text{Fe}/\text{H}]$ relation of Israelian *et al.* (1998) and Boesgaard *et al.* (1998) since the required oxygen yields are unrealistic. Thus the subject is controversial.

Concerning lithium, a compilation of the data is shown in Lemoine *et al.* (1997) and Molaro *et al.* (1997). The Spite plateau extends up to $[\text{Fe}/\text{H}] = -1.3$. Beyond, Li/H is strongly increasing until its solar value of $2 \cdot 10^{-9}$.

A stringent constraint to any theory of Li evolution is avoiding to cross the Spite's plateau below $[\text{Fe}/\text{H}] = -1$. Accordingly, the Li/Be production ratio should be less than about 100.

Recent measurements of ${}^6\text{Li}$ have been made successfully in two halo stars, HD84937 and BD +26 3578 at about $[\text{Fe}/\text{H}] = -2.3$ (Hobbs and Thorburn 1997, Cayrel *et al.* 1998, Smith *et al.* 1998), yielding ${}^6\text{Li}/{}^7\text{Li}$ about 0.05. The great interest of ${}^6\text{Li}$, besides of being an indicator of stellar destruction (Pinsonneault *et al.* 1998, Chaboyer 1998, Cayrel *et al.* 1999, Vauclair and Charbonnel 1998) is to represent a pure spallation product as ${}^9\text{Be}$.

Lithium-6 has different sources, a secondary one and two primary ones: i) fast (p, α) on CNO at rest (this secondary process related to GCRs should not be efficient in the early galaxy), ii) fast CO on H, He (primary) and iii) specifically $\alpha + \alpha$ at low energy (primary). The non thermal fusion reaction $\alpha + \alpha$ produces almost equal amounts of ${}^6\text{Li}$ and ${}^7\text{Li}$ at low energy and the cross section above 100 MeV is specially low (Read and Viola 1984). The second and third processes are associated to LEC. Consequently, this LEC is specially fertile in lithium isotopes.

Preliminary estimates of the ${}^6\text{Li}/{}^9\text{Be}$ ratio have been performed for few Pop II stars but with a large uncertainty (20 - 80). This range of values is much higher than the ${}^6\text{Li}/{}^9\text{Be}$ ratio generated by the present GCR (6). This could indicate that this ratio is varying all along the galactic evolution (Vangioni-Flam *et al.* 1998b, Fields and Olive 1998).

To summarize, we can give six observational constraints on LiBeB evolution :

1. Be and B proportional to Fe
2. $\text{Li}/\text{Be} < 100$ up to $[\text{Fe}/\text{H}] = -1$

3. B/Be = 10-30
4. $^{11}\text{B}/^{10}\text{B} = 4$ at solar birth
5. $^7\text{Li}/^6\text{Li} = 12.5$ at solar birth
6. $^6\text{Li}/^7\text{Li} = 0.05$ and $^6\text{Li}/^9\text{Be} = 20$ to 80 (to be confirmed) at [Fe/H] about -2.3

We recall that the observational O - Fe relation is central to the interpretation since specifically the production of Be is related to O. Most of the observers find oxygen proportional to Fe.

2. Basic physical parameters of non thermal nucleosynthesis of LiBeB

Four parameters are influential to the spallative production of light elements: the reaction cross sections, the energy spectrum of fast nuclei, the composition of the beam and the composition of the target.

Cross sections are well measured (Read and Viola 1984) and have been updated recently by Ramaty *et al.* (1997).

The adopted spectra are of two kinds:

1. GCR: $N(E)dE = kE^{-2.7}$ above a few GeV/n with a flatening below (e.g. Lemoine *et al.* 1998).
2. LEC : Shock wave acceleration with a cut at E_0 of the form $N(E) dE = kE^{-1.5} \exp(-E/E_0) dE$ (Ramaty *et al.* 1996), propagated in the ISM.

The source composition of GCR is well determined (e.g. Du Vernois 1996). It is p and α rich (H/O = 200, He/O = 20) contrary to the possible source composition of the LEC. The most obvious contributors to LEC are supernovae, Wolf-Rayet and mass losing stars (Cassé *et al.* 1995, Ramaty *et al.* 1996, 1997, 1998). It is worth noting that in the early galaxy, supernovae play a leading role since at very low metallicity the stellar winds are insignificant. Table 1 shows a sample of compositions used by different authors : solar system (SS) for comparison (Ramaty *et al.* 1996 from Anders and Grevesse 1989), cosmic ray source (CRS) (Ramaty *et al.* 1996 from Mewaldt 1983), wind of massive stars (W40) (Parizot *et al.* 1997 from Meynet *et al.* 1994), composition of grain products (GR) (Ramaty *et al.* 1996 and Lingenfelter *et al.* 1998), 40 Mo supernova at $Z = 10^{-4} Z_\odot$ (Parizot *et al.* 1999 from Woosley and Weaver 1995) , 35 Mo supernova of solar metallicity (Ramaty *et al.* 1996 from Weaver and Woosley 1993). The two supernovae, though at different metallicities, (SN40 and SN35), give similar yields due to the fact that metallicity dependent mass loss has not been taken into account. Resulting elemental and isotopic ratios (B/Be, $^{11}\text{B}/^{10}\text{B}$, $^6\text{Li}/^9\text{Be}$) for different compositions and E_0 can be found in Ramaty *et al.* (1996) and Vangioni-Flam *et al.* (1997).

TABLE 1 : SOURCE COMPOSITION

Element	SS	CRS	W40	GR	SN40(low Z)	SN35(Zo)
H	1200	220	80	2	37	27
He	120	22	25	0	8.8	7.6
C	0.47	0.87	1.6	0.3	0.09	0.08
N	0.13	0.04	-	0.03	-	-
O	1	1	1	1	1	1

Note that N is insignificant and that Type II supernovae are O-rich whereas the winds of massive stars are C and He-rich. These abundance differences are important since the highest $^{11}\text{B}/^{10}\text{B}$ ratios are produced by C-rich beams through $^{12}\text{C}(\text{p},\text{x})^{11}\text{B}$ and the highest $^6\text{Li}/^9\text{Be}$ ratios are produced by He and O rich compositions.

The fourth parameter, i.e., the composition of the target (ISM) varies from the birth of the galaxy up to now. The extensive study of Ramaty *et al.* (1996) and Vangioni-Flam *et al.* (1997) shows that there are only slight differences in the results when the ISM metallicity is varied between 0 (early galaxy) and Zo (now), except perhaps concerning the $^{11}\text{B}/^{10}\text{B}$ ratio.

3. LiBeB production mechanisms and galactic evolution

Analyzing all the physical parameters discussed above, two main LiBeB producers emerge, the first one is the standard GCR (spectrum 1) in which fast p, α nuclei interact with CNO in the ISM. This process seems unable to produce sufficient amounts of LiBeB at the level observed in the halo stars (Vangioni-Flam *et al.* 1996). A recent study (Fields and Olive 1998) based on the O - Fe relation derived by Israelian *et al.* (1998) and Boesgaard *et al.* (1998) at low metallicity (still controversial) try to fit the observational constraints with a pure standard GCR (secondary production) component, but has problems with the B/Be ratio among other things.

The second one, reverse to the GCR mechanism, invokes fragmentation of CO nuclei in flight by collision with H and He in the ISM. Massive stars are able in principle to furnish freshly synthesized C and O and accelerate them via the shock waves they induce. This mechanism is related to superbubbles (S) through a scenario proposed by Bykov (1995), Parizot (1998) and Vangioni-Flam *et al.* (1998a). Here only the most massive stars (greater than about 60 Mo) contribute due their short lifetime. Originally proposed in relation with the observation of gamma ray line from Orion it remains as a distinctive possibility after the withdrawal of the COMPTEL

results on this molecular complex (Bloemen *et al.* 1998) and the announcement of a possible excess of gamma rays in the 3-7 MeV range from the Vela region. Possible observations in X-rays seem to substantiate this scenario (Tatischeff *et al.* 1998). The observation or non observation of C, O lines at 4.4 and 6.1 MeV and of the Li-Be feature close to 500 keV by the INTEGRAL satellite (Winkler 1997) will be the strongest test of the superbubble hypothesis (Parizot *et al.* 1997).

The acceleration of grain debris in supernovae (scenario G) is another version of a primary mechanism (Lingenfelter *et al.* 1998, Ramaty, these proceedings). The G model leads to quite different predictions concerning the evolution of Be, B and ^6Li only at very low metallicity (Vangioni-Flam *et al.* 1998a,b) because all SN II (10 - 100 Mo) are implied rather than the most massive ones in the S scenario.

Finally, neutrino spallation (Woosley *et al.* 1990, Woosley and Weaver 1995, Vangioni-Flam *et al.* 1996) is helpful to increase the $^{11}\text{B}/^{10}\text{B}$ ratio up to the value observed in meteorites. It is also a primary process since it implies the break up of ^{12}C within supernovae and not in the ISM. However it cannot be the unique mechanism to produce light elements since it does not produce ^9Be . Moreover, it has been shown that its contribution is only marginal (Vangioni-Flam *et al.* 1996).

These different mechanisms are included in a galactic evolutionary model (Vangioni-Flam *et al.* 1996, 1998a,b) to follow the whole evolution of each isotope.

Concerning beryllium and boron, in this context, the main results are the following: the quasi-linearity (Be-B vs Fe) is easily reproduced (fig 2). Standard GCR contribute no more than about 30 per cent to Solar System values. The B/Be ratio is in the range 10-30 as observed. The value 30 leaves enough room for neutrino spallation to reach $^{11}\text{B}/^{10}\text{B} = 4$ at solar birth.

The $^6\text{Li}/\text{H}$ ratio can be explained in the framework of the same superbubble model (Vangioni-Flam *et al.* 1998b, Cayrel *et al.* 1998) this without piercing the Spite plateau (fig 3). In this figure, showing the evolution of $^6\text{Li}/\text{H}$ vs $[\text{Fe}/\text{H}]$, it can be seen that GCR is overwhelmed by LEC. The decrease of the $^6\text{Li}/^9\text{Be}$ ratio could be explained in terms of the variation of the composition of superbubbles in the course of the galactic evolution, being O rich at start due to SNII and becoming more and more C rich due to the increasing contribution of mass losing stars (Vangioni Flam *et al.* 1998b). Moreover, the evolutionary curve of ^6Li crosses the halo observations (fig 3) meaning that ^6Li is almost essentially intact in the envelope of stars in which it is measured. ^7Li in turn, more tightly bound than ^6Li , is even less destroyed, thus the mean value of the Spite plateau reflects nicely the Big Bang ^7Li abundance. This reinforce the use of ^7Li as a cosmological

baryometer.




Figure 3. Evolution of Lithium vs $[\text{Fe}/\text{H}]$

4. CONCLUSION

Recent LiBeB observations indicate that a primary component is probably at work in the early Galaxy, presumably related to core collapse supernovae. Promising scenarios have been presented implying respectively the acceleration of freshly synthesized C and O in superbubbles and/or grains debris around supernovae (see also Ramaty in these proceedings).

New ${}^6\text{Li}$ observations put strong constraints on the composition and spectrum of an early population of fast particles. The superbubble model is also able to reproduce the ${}^6\text{Li}$ observation, until the local meteoritic value (fig 3). A low energy component originally O rich and becoming progressively C rich due to the strengthening of stellar winds at increasing metallicities is required to explain the high ${}^6\text{Li}/{}^9\text{Be}$ observed in a few halo stars with respect to the one measured in meteorites. But a definitive conclusion should wait confirmation.

Essentially no destruction of ${}^6\text{Li}$ and ${}^7\text{Li}$ is implied by the evolutionary curve of ${}^6\text{Li}$. As a consequence, ${}^7\text{Li}$ is a good baryonic density indicator for cosmology. The needs for the future are the following:

On the theoretical side it would be necessary:

- i) to check NLTE corrections on B abundances since two bothering points remain at very low Z .
- ii) to develop and refine SN II models, specially at very low Z and high mass, $M \gtrsim 60$ Mo.

On the observational side, it would be desirable to get measurements of ${}^6\text{Li}$, ${}^7\text{Li}$, ${}^9\text{Be}$, B, O, Fe in the same halo stars and to get ${}^{11}\text{B}/{}^{10}\text{B}$ ratios in

various stars. A first step in this goal has been accomplished by Rebull *et al.* (1998).

Finally, the observation of C,O and Li-Be gamma-ray lines are important objectives of the INTEGRAL satellite which will open up an European era in gamma-ray astronomy (Parizot *et al.* 1997, Cassé *et al.* 1998).

5. Acknowledgements

We have been very much honored to participate to the celebration of the scientific work of Giusa and Roger Cayrel, who are an example for every researcher in Astronomy. Moreover, we thank warmly Monique and Francois Spite for the excellent organization of the meeting. This work was supported in part by the PICS 319, CNRS.

References

- Anders, E. and Grevesse, N. 1989, *Geochim. Acta* 54, 197
 Bloemen, H., 1998, in "The Extreme Universe", Third INTEGRAL Workshop, Taormina
 ESA proceedings, to appear
 Boesgaard, A.M. and King, J.R. 1993, *AJ*, 106, 2309
 Boesgaard, A. *et al.* 1998, preprint
 Burbidge, E.M., Burbidge, G.R., Fowler, W.A. and Hoyle, F. 1957, *Rev.Mod.Phys.* 29, 547
 Bykov, A.M., 1995, *Space Sci.Rev.* 74, 397
 Cassé, M., Lehoucq, R. and Vangioni-Flam, E. 1995, *Nature*, 373, 318
 Cassé, M., Vangioni-Flam, E. and Paul, J., 1998, *International Nuclear Physics Conference*, Paris, Nuclear Physics A, to appear
 Cayrel, R., 1996, *A.A. Rev.* 7, 217
 Cayrel, R., Spite, M., Spite, F., Vangioni-Flam, E., Cassé, M. and Audouze, J. 1998, *AA* accepted
 Cayrel, R., Lebreton, Y., Morel, P., 1999, this proceedings
 Chaboyer, B. 1998, submitted
 Duncan, D.K., Lambert, D. and Lemke, M 1992, *ApJ* 401, 584
 Duncan, D.K. et al 1997, *ApJ*, 488, 333
 Du Vernois, M.A., 1996, in "Cosmic Abundances", Edts ASP Conferences series, Vol 99, p.385
 Fields, B. and Olive K., 1998, *ApJ* submitted
 García-López, R.J. *et al.* 1998, *ApJ*, in press
 Gilmore G., Gustafsson B., Edvardsson B. and Nissen P.E., 1992, *Nature* 357, 379
 Hobbs, L.M. and Thorburn, J.A. 1997, *ApJ* 491, 772
 Israelian, G., García-López, R.J. and Rebolo, R. 1998, *ApJ*, to be published
 Kiselman, D. and Carlsson, M 1996, *AA* 311, 681
 Lemoine, M., Schramm, D.N., Truran, J.W. and Copi, C.J., 1997, *ApJ*, 478, 554
 Lemoine, M., Vangioni-Flam, E. and Cassé, M. 1998, *ApJ*, 499, 735
 Lingenfelter, R.E., Ramaty, R. and Kozlovsky, B. 1998, *ApJL*, 500, in press
 Mac Williams, A., 1997, *Ann.Rev.Astron.Astrophys.*, 55, 503
 Meneguzzi, M., Audouze, J. and Reeves, H., 1971, *AA*, 15, 337
 Mewaldt, R., 1983, *Rev. Geophys. Space Phys.*, 21, 295
 Meynet, G. *et al.* , 1994, *AAS*, 103, 97
 Molaro, P., Bonifacio, P., Castelli, F. and Pasquini, L. 1997, *AA* 319, 593

- Parizot, E., Cassé, M., Vangioni-Flam, E. 1997, AA 328, 107
Parizot, E., 1998, AA to appear
Parizot, E. *et al.*, 1999, in preparation
Pinsonneault, M.H., *et al.* 1998, ApJ, submitted
Raisbeck, G. and Yiou, F., 1971, Phys.Rev. A, 4, 1848
Raisbeck, G. and Yiou, F. 1975, in Spallation Nuclear Reactions and their Applications, Ed B.S.P. Shen and M. Merker (Dordrecht, Reidel), p.83
Ramaty, R., Kozlovsky, B. and Lingenfelter, R.E. 1996, ApJ 456, 525
Ramaty, R., Kozlovsky, B., Lingenfelter, R.E. and Reeves, H. 1997, ApJ, 488, 730
Ramaty, R., Kozlovsky, B. and Lingenfelter, R.E. 1998, Physics Today, 51, no 4, 30
Read, S.M. and Viola, V.E., 1984, Atomic Data and Nuclear Data Tables, 31, 359
Reeves, H., 1994 Rev. of Mod. Phys., Vol 66, No 1, p. 193
Rebolo, R., Molaro, P., Beckman J.E. 1988, AA, 192, 192
Rebull, L.M. *et al.*, 1998, preprint
Ryan S., Norris I., Bessel M. and Deliyannis C. 1994, Astrophys. J. 388, 184
Ryan, S.G., Norris, J.E. and Beers, T.C. 1996, ApJ, 471, 254
Smith, V.V., Lambert, D.L. and Nissen, P.E. 1993, ApJ, 408, 262
Smith, V.V., Lambert, D.L. and Nissen, P.E., 1998, ApJ, 506, 405
Tatischeff, V., Ramaty, R. and Kozlovsky, B., 1998, ApJ to be published
Thielemann, F.K., Nomoto, K. and Hashimoto, M., 1996, ApJ, 460, 108
Vangioni-Flam E., Cassé, M., Olive, K. and Fields B.D 1996, ApJ 468, 199
Vangioni-Flam, E., Cassé, M. and Ramaty, R. 1997, in Proceedings of the 2nd INTEGRAL Workshop, "The Transparent Universe", Saint Malo, France, ESA, SP.382 p. 123
Vangioni-Flam, E., Ramaty, R., Olive, K. and Cassé, M. 1998a, AA 337, 714
Vangioni-Flam, E., Cassé, M., Cayrel, R., Audouze, J., Spite, M., Spite, F. 1998b, New Astronomy, submitted
Vauclair, S. and Charbonnel, C. 1998, AA, ApJ, 502, 372
Weaver, T.A. and Woosley, S.E., 1993, Phys. Repts, 227, 65
Winkler, C., 1997, in "The Transparent Universe", 2nd INTEGRAL Workshop, Saint Malo, Edts, ESA, SP-382, p. 573
Woosley, S.E., Hartmann, D., Hoffman, R., Haxton, W., 1990, ApJ, 356, 272
Woosley, S.E. and Weaver, T.A. 1995, Ap.JS101, 181